arXiv:0910.4075v1 [math.DG] 21 Oct 2009

Universal bounds on eigenvalues of the buckling problem on spherical domains *

Guangyue Huang, Xingxiao Li †, Linfen Cao
Department of Mathematics, Henan Normal University
Xinxiang 453007, Henan, P.R. China

Abstract. In this paper we study the eigenvalues of buckling problem on domains in a unit sphere. By introducing a new parameter and using Cauchy inequality, we optimize the inequality obtained by Wang and Xia in [12].

Keywords: eigenvalue, universal bounds, buckling problem.
Mathematics Subject Classification: Primary 35P15, Secondary 58G25.

1 Introduction

Let Ω be a connected bounded domain in an $n(\geq 2)$-dimensional Euclidean space $\mathbb{R}^n$ and ν be the unit outward normal vector field of $\partial\Omega$. The well-known eigenvalue problem

$$\begin{cases} \Delta^2 u = \Lambda(-\Delta)u & \text{in } \Omega, \\ u = \frac{\partial u}{\partial \nu} = 0 & \text{on } \partial\Omega \end{cases} \tag{1.1}$$

is called a *buckling problem*, which is used to describe the critical buckling load of a clamped plate subjected to a uniform compressive force around its boundary.

Let

$$0 < \Lambda_1 \leq \Lambda_2 \leq \Lambda_3 \leq \cdots$$

denote the successive eigenvalues for (1.1), where each eigenvalue is repeated according to its multiplicity. In 1956, Payne-Pólya-Weinberger [11] proved

$$\Lambda_2 \leq 3\Lambda_1 \quad \text{for } \Omega \in \mathbb{R}^2. \tag{1.2}$$

Following the method of Payne-Pólya-Weinberger in [11], it reads that for $\Omega \in \mathbb{R}^n$ (for the generalization of (1.2) to n dimensions, see [2]):

$$\Lambda_2 \leq \left(1 + \frac{8}{n+2}\right)\Lambda_1.$$

---

[*] This research is supported by NSFC of China (No. 10671181), Project of Henan Provincial department of Sciences and Technology (No. 092300410143), and NSF of Henan Provincial Education department (No. 2009A110010).

[†] The corresponding author. Email: xxl@henannu.edu.cn

Subsequently, in 1984, Hile and Yeh [7] improved the above inequality as follows:

$$\Lambda_2 \leq \frac{n^2+8n+20}{(n+2)^2}\Lambda_1.$$

In 1998, Ashbaugh [1] obtained

$$\sum_{i=1}^{n} \Lambda_{i+1} \leq (n+4)\Lambda_1.$$

In a recent survey paper, answering a question of Ashbaugh in [1], Cheng-Yang [4] proved the following universal inequalities on eigenvalues for the eigenvalue problem (1.1):

$$\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2 \leq \frac{4(n+2)}{n^2}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\Lambda_i.$$

It has become clear that many similar inequalities on eigenvalues of Laplacian of Payne-Pólya-Weinberger rely on facts involving operators, their commutators, and traces. For the related research and improvement in this direction, see [3,5,6,8–10,13] and the references therein.

Let $x_1, x_2, \ldots, x_{n+1}$ be the standard Euclidean coordinate functions of $\mathbb{R}^{n+1}$. Then the unit sphere is defined by

$$\mathbb{S}^n = \left\{ (x_1, x_2, \ldots, x_{n+1}) \in \mathbb{R}^{n+1} \ ; \ \sum_{\alpha=1}^{n+1} x_\alpha^2 = 1 \right\}.$$

In 2007, Wang and Xia [12] considered the buckling problem on domains in a unit sphere $\mathbb{S}^n$ and obtained the following result:

Theorem 1.1. [12] *Let Λ_i be the i^{th} eigenvalue of the following eigenvalue problem:*

$$\Delta^2 u = \Lambda(-\Delta)u \ \text{ in } \Omega, \qquad u = \frac{\partial u}{\partial \nu} = 0 \ \text{ on } \partial\Omega,$$

where Ω is a connected domain in a unit sphere $\mathbb{S}^n (n \geq 2)$ with smooth boundary $\partial\Omega$ and ν is the unit outward normal vector field of $\partial\Omega$. Then for any $\delta > 0$,

$$\begin{aligned} 2\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2 \leq & \sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\delta\Lambda_i + \frac{\delta^2(\Lambda_i-(n-2))}{4(\delta\Lambda_i+n-2)}\right) \\ & + \frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i + \frac{(n-2)^2}{4}\right). \end{aligned} \tag{1.3}$$

We remark that the right hand side of inequality (1.3) depends on δ. In the current paper, by introducing a new parameter and using Cauchy inequality, we obtain a stronger inequality than (1.3) which is independent of δ, and derive an inequality of the type of Yang (see inequality (1.5)). Our main results are stated as follows:

Theorem 1.2. *Let Λ_i be the i^{th} eigenvalue of the following eigenvalue problem:*

$$\Delta^2 u = \Lambda(-\Delta)u \ \text{ in } \Omega, \qquad u = \frac{\partial u}{\partial \nu} = 0 \ \text{ on } \partial\Omega,$$

where Ω is a connected domain in a unit sphere $\mathbb{S}^n(n \geq 2)$ with smooth boundary $\partial\Omega$ and ν is the unit outward normal vector field of $\partial\Omega$. Then

$$\begin{aligned}
&\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(2+\frac{n-2}{\Lambda_i-(n-2)}\right)\\
\leq& 2\left\{\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\Lambda_i-\frac{n-2}{\Lambda_i-(n-2)}\right)\right\}^{1/2}\\
&\times\left\{\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right)\right\}^{1/2}. \qquad (1.4)
\end{aligned}$$

Corollary 1.3. *Under the assumptions of Theorem 1.2,*

$$\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2 \leq \sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i-\frac{n-2}{\Lambda_i-(n-2)}\right)\left(\Lambda_i+\frac{(n-2)^2}{4}\right), \qquad (1.5)$$

$$\Lambda_{k+1} \leq S_{k+1}+\sqrt{S_{k+1}^2-T_{k+1}}, \qquad (1.6)$$

$$\Lambda_{k+1}-\Lambda_k \leq 2\sqrt{S_{k+1}^2-T_{k+1}}, \qquad (1.7)$$

where

$$S_{k+1}=\frac{1}{k}\sum_{i=1}^{k}\Lambda_i+\frac{1}{2k}\sum_{i=1}^{k}\left(\Lambda_i-\frac{n-2}{\Lambda_i-(n-2)}\right)\left(\Lambda_i+\frac{(n-2)^2}{4}\right), \qquad (1.8)$$

$$T_{k+1}=\frac{1}{k}\sum_{i=1}^{k}\Lambda_i^2+\frac{1}{k}\sum_{i=1}^{k}\Lambda_i\left(\Lambda_i-\frac{n-2}{\Lambda_i-(n-2)}\right)\left(\Lambda_i+\frac{(n-2)^2}{4}\right). \qquad (1.9)$$

Remark 1.4. The inequality (1.4) is equivalent to

$$\begin{aligned}
2\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2 \leq& -\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\frac{n-2}{\Lambda_i-(n-2)}\\
&+2\left\{\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\Lambda_i-\frac{n-2}{\Lambda_i-(n-2)}\right)\right\}^{1/2}\\
&\times\left\{\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right)\right\}^{1/2}.
\end{aligned}$$

From the inequality (2.15) in Section 2 and making use of Cauchy inequality, the following inequality can be deduced:

$$
\begin{aligned}
&-\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\frac{n-2}{\Lambda_i-(n-2)}\\
&\quad+2\left\{\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\Lambda_i-\frac{n-2}{\Lambda_i-(n-2)}\right)\right\}^{1/2}\\
&\quad\times\left\{\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right)\right\}^{1/2}\\
&\le-\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\frac{n-2}{\Lambda_i-(n-2)}\\
&\quad+\delta\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\Lambda_i-\frac{n-2}{\Lambda_i-(n-2)}\right)\\
&\quad+\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right)\\
&=\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\delta\Lambda_i-\frac{(\delta+1)(n-2)}{\Lambda_i-(n-2)}\right)\\
&\quad+\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right)\\
&\le\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\delta\Lambda_i+\frac{\delta^2(\Lambda_i-(n-2))}{4(\delta\Lambda_i+n-2)}\right)\\
&\quad+\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right),
\end{aligned}
$$

which shows that the inequality (1.4) is sharper than inequality (1.3). Therefore, Theorem 1.2 improves Theorem 1.1.

2 Proof of Theorem

By the method of constructing trial functions introduced by Cheng and Yang in [4], for any $\alpha=1,\dots,n+1$ and each $i=1,\dots,k$, the vector-valued functions $x_\alpha\nabla u_i$ can be decomposed as

$$x_\alpha\nabla u_i=\nabla h_{\alpha i}+W_{\alpha i}, \tag{2.1}$$

where $h_{\alpha i}\in H^2_{2,D}(\Omega)$, $\nabla h_{\alpha i}$ is the projection of $x_\alpha\nabla u_i$ in $\mathbf{H}^2_{1,D}(\Omega)$ and $W_{\alpha i}\perp\mathbf{H}^2_{1,D}(\Omega)$ (for the definitions of $H^2_{2,D}(\Omega)$ and $\mathbf{H}^2_{1,D}(\Omega)$, we refer to [12]). Hence,

$$W_{\alpha i}|_{\partial\Omega}=0$$

and

$$\int_\Omega \langle W_{\alpha i}, \nabla u\rangle = 0, \qquad \text{for any } u \in H^2_{2,D}(\Omega).$$

Define $\|f\|^2 = \int_\Omega |f|^2$. Then (see inequalities (2.19) and (2.40) in [12])

$$\|x_\alpha \nabla u_i\|^2 = \|\nabla h_{\alpha i}\|^2 + \|W_{\alpha i}\|^2 \tag{2.2}$$

and for any $\delta > 0$,

$$\begin{aligned}\sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)^2 r_{\alpha i} \leq &\sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)^2 \Big(\delta p_{\alpha i} + (\delta\Lambda_i + n - 2)\|W_{\alpha i}\|^2 \\ &+ \delta\|\langle \nabla x_\alpha, \nabla u_i\rangle\|^2\Big) + \frac{1}{\delta}\sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)\|Z_{\alpha i}\|^2,\end{aligned} \tag{2.3}$$

where

$$\begin{aligned} r_{\alpha i} =& 2\|\langle \nabla x_\alpha, \nabla u_i\rangle\|^2 + \int_\Omega \langle \nabla x_\alpha^2, \Delta u_i \nabla u_i\rangle + (n-2)\|x_\alpha \nabla u_i\|^2, \\ p_{\alpha i} =& \int_\Omega \langle \nabla x_\alpha^2, u_i \nabla(\Delta u_i) + \Lambda_i u_i \nabla u_i\rangle, \\ Z_{\alpha i} =& \nabla\langle \nabla x_\alpha, \nabla u_i\rangle - \frac{n-2}{2} x_\alpha \nabla u_i. \end{aligned}$$

Next, we are to optimize the inequality (2.41) in [12]. Let C be a positive constant. Then it follows from Cauchy inequality that

$$\begin{aligned} C\|\langle \nabla x_\alpha, \nabla u_i\rangle\|^2 =& C\int_\Omega \langle \nabla x_\alpha, \nabla u_i\rangle^2 \\ =& -C\int_\Omega x_\alpha \mathrm{div}(\langle \nabla x_\alpha, \nabla u_i\rangle \nabla u_i) \\ =& -C\int_\Omega \langle x_\alpha \nabla u_i, \nabla\langle \nabla x_\alpha, \nabla u_i\rangle\rangle - C\int_\Omega \langle \nabla x_\alpha, \nabla u_i\rangle x_\alpha \Delta u_i \\ =& -C\int_\Omega \langle \nabla h_{\alpha i}, \nabla\langle \nabla x_\alpha, \nabla u_i\rangle\rangle - \frac{C}{2}\int_\Omega \langle \nabla x_\alpha^2, \nabla u_i\rangle \Delta u_i \\ \leq& (\delta\Lambda_i + n - 2)\|\nabla h_{\alpha i}\|^2 + \frac{C^2}{4(\delta\Lambda_i + n - 2)}\|\nabla\langle \nabla x_\alpha, \nabla u_i\rangle\|^2 \\ & - \frac{C}{2}\int_\Omega \langle \nabla x_\alpha^2, \nabla u_i\rangle \Delta u_i, \end{aligned} \tag{2.4}$$

where div(Z) denotes the divergence of Z. Applying (2.4) to (2.3) yields

$$\begin{aligned}
\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2 r_{\alpha i} \le{}& \sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\Big(\delta p_{\alpha i}+(\delta\Lambda_i+n-2)\|W_{\alpha i}\|^2 \\
&+(\delta-C)\|\langle\nabla x_\alpha,\nabla u_i\rangle\|^2+C\|\langle\nabla x_\alpha,\nabla u_i\rangle\|^2\Big) \\
&+\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\|Z_{\alpha i}\|^2 \\
\le{}& \sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\Big(\delta p_{\alpha i}+(\delta\Lambda_i+n-2)(\|W_{\alpha i}\|^2+\|\nabla h_{\alpha i}\|^2) \\
&+(\delta-C)\|\langle\nabla x_\alpha,\nabla u_i\rangle\|^2+\frac{C^2}{4(\delta\Lambda_i+n-2)}\|\nabla\langle\nabla x_\alpha,\nabla u_i\rangle\|^2 \\
&-\frac{C}{2}\int_\Omega\langle\nabla x_\alpha^2,\nabla u_i\rangle\Delta u_i\Big)+\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\|Z_{\alpha i}\|^2 \\
={}& \sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\Big(\delta p_{\alpha i}+(\delta\Lambda_i+n-2)\|x_\alpha\nabla u_i\|^2 \\
&+(\delta-C)\|\langle\nabla x_\alpha,\nabla u_i\rangle\|^2+\frac{C^2}{4(\delta\Lambda_i+n-2)}\|\nabla\langle\nabla x_\alpha,\nabla u_i\rangle\|^2 \\
&-\frac{C}{2}\int_\Omega\langle\nabla x_\alpha^2,\nabla u_i\rangle\Delta u_i\Big)+\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\|Z_{\alpha i}\|^2, \qquad (2.5)
\end{aligned}$$

where in the last step in (2.5) we have used equality (2.2). A direct calculation yields (see (2.44), (2.45), (2.46) and (2.47) in [12])

$$\sum_{\alpha=1}^{n+1} r_{\alpha i}=n, \qquad (2.6)$$

$$\sum_{\alpha=1}^{n+1} p_{\alpha i}=0, \qquad (2.7)$$

$$\sum_{\alpha=1}^{n+1}\|x_\alpha\nabla u_i\|^2=\sum_{\alpha=1}^{n+1}\|\langle\nabla x_\alpha,\nabla u_i\rangle\|^2=1, \qquad (2.8)$$

$$\sum_{\alpha=1}^{n+1}\|\nabla\langle\nabla x_\alpha,\nabla u_i\rangle\|^2=\Lambda_i-(n-2) \qquad (2.9)$$

and

$$\sum_{\alpha=1}^{n+1}\|Z_{\alpha i}\|^2=\Lambda_i+\frac{(n-2)^2}{4}. \qquad (2.10)$$

Therefore, summing up (2.5) over α from 1 to $n+1$, one gets

$$\begin{aligned} n\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2 \leq & \sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\Big(\delta\Lambda_i+n-2+(\delta-C) \\ & +\frac{C^2}{4(\delta\Lambda_i+n-2)}(\Lambda_i-(n-2))\Big) \\ & +\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right). \end{aligned}$$

That is,

$$\begin{aligned} 2\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2 \leq & \sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\delta\Lambda_i+(\delta-C)\right. \\ & \left.+\frac{C^2}{4(\delta\Lambda_i+n-2)}(\Lambda_i-(n-2))\right) \\ & +\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right). \end{aligned} \tag{2.11}$$

Taking

$$C=\frac{2(\delta\Lambda_i+n-2)}{\Lambda_i-(n-2)}$$

in (2.11) yields

$$\begin{aligned} 2\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2 \leq & \sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\delta\Lambda_i-\frac{(\delta+1)(n-2)}{\Lambda_i-(n-2)}\right) \\ & +\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right), \end{aligned} \tag{2.12}$$

and hence

$$\begin{aligned} & \sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(2+\frac{n-2}{\Lambda_i-(n-2)}\right) \\ \leq & \delta\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)^2\left(\Lambda_i-\frac{n-2}{\Lambda_i-(n-2)}\right) \\ & +\frac{1}{\delta}\sum_{i=1}^{k}(\Lambda_{k+1}-\Lambda_i)\left(\Lambda_i+\frac{(n-2)^2}{4}\right). \end{aligned} \tag{2.13}$$

To complete the proof of Theorem 1.2, we need the following lemma:

Lemma 2.1. *Let Ω be a connected bounded domain in $\mathbb{S}^n$. Then*

$$\Lambda_1 \geq n. \tag{2.14}$$

Proof. Let Ω_1, Ω_2 be two connected bounded domains in $\mathbb{S}^n$ and $\Omega_1 \subset \Omega_2$. Let $u_1(\Omega_1)$ be the eigenfunction corresponding to $\Lambda_1(\Omega_1)$. Then the function defined by

$$\widetilde{u_1} = \begin{cases} u_1 & \text{in } \Omega_1, \\ 0 & \text{in } \Omega_2 - \Omega_1 \end{cases}$$

is a eigenfunction corresponding to $\Lambda_1(\Omega_2)$. Hence, $\Lambda_1(\Omega_1) \geq \Lambda_1(\Omega_2)$. Denote by λ_1 the first eigenvalue of Laplacian. It is easy to see $\Lambda_1(\mathbb{S}^n) = \lambda_1(\mathbb{S}^n) = n$ because there are no boundary conditions in this case. It follows that $\Lambda_1(\Omega) \geq \Lambda_1(\mathbb{S}^n) = n$ by setting $\Omega = \Omega_1$ and $\Omega_2 = \mathbb{S}^n$. This completes the proof of Lemma 2.1. □

Inequality (2.14) shows that for any i,

$$\Lambda_i - \frac{n-2}{\Lambda_i - (n-2)} > 0. \tag{2.15}$$

Minimizing the right hand side of (2.13) as a function of δ by choosing

$$\delta = \left(\frac{\sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)\left(\Lambda_i + \frac{(n-2)^2}{4}\right)}{\sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)^2 \left(\Lambda_i - \frac{n-2}{\Lambda_i - (n-2)}\right)} \right)^{1/2},$$

we obtain the inequality (1.4), completing the proof of Theorem 1.2.

Proof of Corollary 1.3. It is easy to see from (1.4) that

$$\sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)^2 \leq \left\{ \sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)^2 \left(\Lambda_i - \frac{n-2}{\Lambda_i - (n-2)}\right) \right\}^{1/2} \times \left\{ \sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)\left(\Lambda_i + \frac{(n-2)^2}{4}\right) \right\}^{1/2}. \tag{2.16}$$

One can check by induction that

$$\begin{aligned} &\left\{ \sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)^2 \left(\Lambda_i - \frac{n-2}{\Lambda_i - (n-2)}\right) \right\} \left\{ \sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)\left(\Lambda_i + \frac{(n-2)^2}{4}\right) \right\} \\ &\leq \left\{ \sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)^2 \right\} \left\{ \sum_{i=1}^k (\Lambda_{k+1} - \Lambda_i)\left(\Lambda_i - \frac{n-2}{\Lambda_i - (n-2)}\right)\left(\Lambda_i + \frac{(n-2)^2}{4}\right) \right\}, \end{aligned}$$

which together with inequality (2.16) yields inequality (1.5).

Solving the quadratic polynomial of Λ_{k+1} in (1.5), we obtain inequality (1.6).

Replacing $k+1$ with k in (1.5), we obtain

$$\sum_{i=1}^{k-1} (\Lambda_k - \Lambda_i)^2 \leq \sum_{i=1}^{k-1} (\Lambda_k - \Lambda_i)\left(\Lambda_i - \frac{n-2}{\Lambda_i - (n-2)}\right)\left(\Lambda_i + \frac{(n-2)^2}{4}\right).$$

Therefore,

$$\sum_{i=1}^{k} (\Lambda_k - \Lambda_i)^2 \leq \sum_{i=1}^{k} (\Lambda_k - \Lambda_i)\left(\Lambda_i - \frac{n-2}{\Lambda_i - (n-2)}\right)\left(\Lambda_i + \frac{(n-2)^2}{4}\right).$$

Solving this inequality of quadratic polynomial for Λ_k, we infer

$$\Lambda_k \geq S_{k+1} - \sqrt{S_{k+1}^2 - T_{k+1}}, \tag{2.17}$$

where S_{k+1}, T_{k+1} are given by (1.8) and (1.9) respectively. Therefore, the inequality (1.7) follows from (2.17) and (1.6). Then Corollary 1.3 is completed.

References

[1] M.S. Ashbaugh, Isoperimetric and universal inequalities for eigenvalues, in Spectral theory and geometry (Edinburgh, 1998) E. B. Davies, and Yu Safarov eds., London Math. Soc. Lecture Notes vol. 273, Cambridge Univ. Press, Cambridge, 1999. pp. 95-139.

[2] M.S. Ashbaugh, On universal inequalities for the low eigenvalues of the buckling problem. (English summary) Partial differential equations and inverse problems, 13-31, Contemp. Math., 362, Amer. Math. Soc., Providence, RI, 2004.

[3] Q.M. Cheng, H.C. Yang, Inequalities for eigenvalues of a clamped plate problem, Trans. Amer. Math. Soc. 358 (2006) 2625-2635.

[4] Q.M. Cheng, H.C. Yang, Universal bounds for eigenvalues of a buckling problem, Commun. Math. Phys. 262 (2006) 663-675.

[5] A. EI Soufi, E.M. Harrell and S. Ilias, Universal inequalities for the eigenvalues of Laplace and Schrödinger operators on submanifolds, Trans. Amer. Math. Soc. 361 (2009) 2337-2350.

[6] E.M. Harrell, J. Stubble, On trace identities and universal eigenvalue estimates for some partial differential operators, Trans. Amer. Math. Soc. 349 (1997) 1797-1809.

[7] G.N. Hile, R.Z. Yeh, Inequalities for eigenvalues of the biharmonic operator, Pacific J. Math. 112 (1984) 115-133.

[8] G.Y. Huang, X.X. Li and R.W. Xu, Extrinsic estimates for the eigenvalues of Schrödinger operator, to appear in Geom. Dedicata.

[9] G.Y. Huang, W.Y. Chen, Inequalities of eigenvalues for the bi-kohn Laplacian on the Heisenberg group, to appear in Acta Math. Sci. Ser. B Engl. Ed. 2009, vol 29(6).

[10] M. Levitin, L. Parnovski, Commutators, spectral trace identities, and universal estimates for eigenvalues, J. Funct. Anal. 192 (2002) 425-445.

[11] L.E. Payne, G. Pólya and H.F. Weinberger, On the ratio of consecutive eigenvalues, J. Math. Phys. 35 (1956), 289-298.

[12] Q.L. Wang, C.Y. Xia, Universal inequalities for eigenvalues of the buckling problem on spherical domains, Commun. Math. Phys. 270 (2007) 759-775.

[13] Q.L. Wang, C.Y. Xia, Universal bounds for eigenvalues of the biharmonic operator on Riemannian manifolds, J. Funct. Anal. 245 (2007) 334-352.